

# One-Sided Confidence Regions for a Multivariate Location Parameter

Michael Vock

Universität Bern  
Institut für mathematische Statistik und Versicherungslehre

ROeS Seminar 2007, Bern, September 11, 2007

Research supported by the Swiss National Science Foundation

# Outline

## Introduction

Direct Derivation of Confidence Regions for  $\vartheta$

Enhanced Confidence Regions for  $\vartheta$

# One-Sided Confidence Regions

- $\mathbf{X}_1, \dots, \mathbf{X}_n$  i. i. d. random vectors in  $\mathbb{R}^p$
- $\mathbf{X}_i \sim P_{\vartheta}$ ,  $\vartheta \in \Theta$  unknown
- $P_{\vartheta}$  (at least directionally) symmetric w. r. t.  $\vartheta$

Problem: Find a  $1 - \alpha$  confidence region for  $\vartheta$  that is

- as strict as possible in specific directions
- possibly unbounded in “irrelevant” directions

(e. g. a cone or an orthant).

## Connection with One-Sided Location Tests

Let  $\varphi_\alpha$  be a non-randomized level  $\alpha$  test for

$$H_0 : \vartheta \in \Theta_0(\gamma) \quad \text{vs.}$$

$$H_1 : \vartheta \in \Theta \setminus \Theta_0(\gamma).$$

(E. g.  $\Theta_0(\gamma) = \gamma + (-\infty, 0]^p$ )

Inversion of  $\varphi_\alpha$

$\Rightarrow \mathcal{C}_{1-\alpha}(\mathbf{X}_1, \dots, \mathbf{X}_n) = \{\gamma : \varphi_\alpha((\mathbf{X}_1, \dots, \mathbf{X}_n), \gamma) = 0\}$ , and

$$P_{\vartheta}(\mathcal{C}_{1-\alpha}(X) \ni \gamma) \geq 1 - \alpha \quad \forall \vartheta \in \Theta_0(\gamma) \quad \forall \gamma \in \Theta.$$

$\mathcal{C}_{1-\alpha}$  is a  $1 - \alpha$  confidence region for the meta-parameter  $\gamma$ .

# Outline

Introduction

**Direct Derivation of Confidence Regions for  $\vartheta$**

Enhanced Confidence Regions for  $\vartheta$

## Direct Derivation of Confidence Regions for $\vartheta$

Assume that  $\gamma \in \Theta_0(\gamma)$ ,  $\forall \gamma \in \Theta$ .

Then

$$P_{\vartheta}(\mathcal{C}_{1-\alpha}(X) \ni \vartheta) \geq 1 - \alpha \quad \forall \vartheta \in \Theta.$$

$\mathcal{C}_{1-\alpha}$  is also a  $1 - \alpha$  confidence region for the location parameter  $\vartheta$ .

## Direct Derivation of Confidence Regions for $\vartheta$

Assume that  $\gamma \in \Theta_0(\gamma)$ ,  $\forall \gamma \in \Theta$ .

Then

$$P_{\vartheta}(\mathcal{C}_{1-\alpha}(X) \ni \vartheta) \geq 1 - \alpha \quad \forall \vartheta \in \Theta.$$

$\mathcal{C}_{1-\alpha}$  is also a  $1 - \alpha$  confidence region for the location parameter  $\vartheta$ .

Problems:

- conservative
- unpleasant shape – to be illustrated...

## Min and Max Tests

### Min Test

Reject  $H_0 : \exists j \in \{1, \dots, p\} : \vartheta_j \leq \gamma_j$  in favor of  $H_1 : \vartheta > \gamma$  at the level  $\alpha$  if and only if

$$\varphi_{j,\alpha}((X_{1j}, \dots, X_{nj}), \gamma_j) = 1, \forall j \in 1, \dots, p.$$


### Bonferroni Max Test

Reject  $H_0 : \vartheta \leq \gamma$  in favor of  $H_1 : \exists j \in \{1, \dots, p\} : \vartheta_j > \gamma_j$  at the level  $\alpha$  if and only if



$$\exists j \in 1, \dots, p : \varphi_{j,\alpha/p}((X_{1j}, \dots, X_{nj}), \gamma_j) = 1.$$

## Example

Two variables of the pulmonary function data by Randles (1989) (slightly modified from Merchant et al., 1975).





# Pulmonary Function Data, Wilcoxon Min Test





$$\mathcal{C}_{1-\alpha}(X) = \mathbf{c}_{1-\alpha}(X) - \Theta_0(\mathbf{0})$$

## P. F. Data, Wilcoxon Bonferroni Max Test



$$\mathcal{C}_{1-\alpha}(X) = \mathbf{c}_{1-\alpha}(X) - \Theta_0(\mathbf{0})$$

## P. F. Data, Sign Test by Larocque/Labarre (2004)



$$\mathcal{C}_{1-\alpha}(X) \approx \mathbf{c}_{1-\alpha}(X) - \Theta_0(\mathbf{0}) \text{ (outside a sufficiently large ball)}$$

## Back to the Drawbacks of the Direct Approach

If confidence regions for  $\gamma$  are directly used as confidence regions for  $\vartheta$ , they are

- usually conservative and
- similar in shape to  $-\Theta_0$ , rather than to  $\Theta_1 = \Theta \setminus \Theta_0$ .

# Outline

Introduction

Direct Derivation of Confidence Regions for  $\vartheta$

**Enhanced Confidence Regions for  $\vartheta$**

# Enhanced Confidence Regions for $\vartheta$ (1)

$$\text{Temptation: } \tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) = \bigcap_{\gamma \notin \mathcal{C}_{1-\alpha}(\mathbf{X})} \Theta_1(\gamma)$$

## Enhanced Confidence Regions for $\vartheta$ (1)

$$\text{Temptation: } \tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) = \bigcap_{\gamma \notin \mathcal{C}_{1-\alpha}(\mathbf{X})} \Theta_1(\gamma)$$

⇒ liberal – multiple testing problem!

Solution: Reduce the set of possible meta-parameters in advance.

## Enhanced Confidence Regions for $\vartheta$ (2)

Let  $\mathcal{C}_{1-\alpha} : \mathcal{X} \rightarrow \mathcal{P}(\mathbb{R}^p)$  be a  $1 - \alpha$  confidence region for  $\gamma$  based on  $(\Theta_0(\gamma))_{\gamma \in \mathbb{R}^p}$ .

Let  $\Theta_0(\gamma) = \gamma + \Theta_0(\mathbf{0})$ ,  $\forall \gamma \in \mathbb{R}^p$ , closed,  $\Theta_1(\gamma) = \mathbb{R}^p \setminus \Theta_0(\gamma)$ .

Assume that  $\Theta_0(\gamma) \subset \Theta_0(\gamma + (\delta, \dots, \delta)^T)$ ,  $\forall \gamma \in \mathbb{R}^p$ ,  $\delta > 0$ .

With  $\gamma_i = (i, \dots, i)^T \in \mathbb{R}^p$ ,  $\forall i \in I = [\ell, \infty)$ , define

$$\tilde{\mathcal{C}}_{1-\alpha}(X) := \bigcap_{i \in I: \gamma_{i'} \notin \mathcal{C}_{1-\alpha}(X) \forall i' \leq i} \Theta_1(\gamma_i).$$


Then

$$P_{\vartheta} \left( \tilde{\mathcal{C}}_{1-\alpha}(X) \ni \vartheta \right) \geq 1 - \alpha \quad \forall \vartheta \in \mathbb{R}^p.$$


## Idea of the Proof

- $(\Theta_0(\gamma_i))_{i \in I}$  is closed under (finite and infinite) intersections.
- Apply the closed testing principle (Marcus, Peritz, and Gabriel, 1976).
- Translate to confidence regions.


# Pulmonary Function Data, Wilcoxon Min Test



# Pulmonary Function Data, Wilcoxon Min Test



# Pulmonary Function Data, Wilcoxon Min Test



## Enhanced Confidence Regions: Properties

- $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) = \mathbf{c}_{1-\alpha}(\mathbf{X}) + \Theta_1(\mathbf{0})$  (by definition)
- $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) \subset \mathcal{C}_{1-\alpha}(\mathbf{X})$  under suitable conditions ( $\Theta_1(\mathbf{0})$  convex cone, translation invariance, and a monotonicity property)

## Enhanced Confidence Regions: Properties

- $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) = \mathbf{c}_{1-\alpha}(\mathbf{X}) + \Theta_1(\mathbf{0})$  (by definition)
- $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X}) \subset \mathcal{C}_{1-\alpha}(\mathbf{X})$  under suitable conditions ( $\Theta_1(\mathbf{0})$  convex cone, translation invariance, and a monotonicity property)

Disadvantage: Restricted set of possible confidence regions (search on the diagonal)

## Summary

- A confidence region  $\mathcal{C}_{1-\alpha}(\mathbf{X})$  obtained by inversion of a test for a composite null hypothesis is for a meta-parameter.
- Even if  $\mathcal{C}_{1-\alpha}(\mathbf{X})$  may also be a confidence region for the parameter  $\vartheta$  itself, it is not very useful.
- The proposed method based on the closed testing principle yields a confidence region  $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X})$  with a more useful shape.
- $\tilde{\mathcal{C}}_{1-\alpha}(\mathbf{X})$  is also less conservative than  $\mathcal{C}_{1-\alpha}(\mathbf{X})$  under suitable conditions.

## References

- Larocque, D., Labarre, M. (2004). A conditionally distribution-free multivariate sign test for one-sided alternatives. *Journal of the American Statistical Association*, **99**, 499–509.
- Marcus, R., Peritz, E., Gabriel, K. R. (1976). On closed testing procedures with special reference to ordered analysis of variance. *Biometrika*, **63**, 655–660.
- Merchant, J. A., Halprin, G. M., Hudson, A. R., Kilburn, K. H., McKenzie, W. N., Hurst, D. J., Bermazohn, P. (1975). Responses to cotton dust. *Archives of Environmental Health*, **30**, 222–229.
- Randles, R. H. (1989). A distribution-free multivariate sign test based on interdirections. *Journal of the American Statistical Association*, **84**, 1045–1050.
- Vock, M. (2007). *Enhanced one-sided confidence regions for a multivariate location parameter*, submitted.