

Actual graft survival

Webtable 7

Associations between OKT3 use and actual graft loss using different model-building strategies

| Parameter | Hazard Ratio | 95 % confidence interval | | p-value |
|-------------------------------|---------------------|---------------------------------|------|----------------|
| <i>Propensity score model</i> | | | | |
| OKT3 vs. ATG usage | 1.55 | 0.91 | 2.63 | 0.110 |
| <i>Automated model</i> | | | | |
| OKT3 vs. ATG usage | 1.27 | 0.75 | 2.15 | 0.382 |
| Year of transplantation | 0.90 | 0.84 | 0.95 | <0.001 |
| <i>Clinical expertise</i> | | | | |
| OKT3 vs. ATG usage | 1.35 | 0.79 | 2.31 | 0.277 |
| Donor age | 1.01 | 1.00 | 1.02 | 0.049 |
| BANFF 2 vs. 1 | 1.34 | 0.95 | 1.88 | 0.098 |
| BANFF 3 vs. 1 | 2.55 | 1.30 | 4.99 | 0.006 |
| HLA mismatch | 1.02 | 0.88 | 1.18 | 0.817 |
| PRA | 1.01 | 0.99 | 1.02 | 0.302 |
| Year of transplantation | 0.88 | 0.83 | 0.94 | <0.001 |