

Webtable 4

Parameters used for propensity score model for actual graft survival

| Parameter | Odds Ratio | 95% confidence interval | | p-value |
|-------------------------------------|-------------------|--------------------------------|-------|----------------|
| Number of bloodpressure medications | 1.17 | 1.04 | 1.31 | 0.008 |
| Peripheral vascular disease | 1.03 | 0.53 | 1.98 | 0.931 |
| Coronary heart disease | 0.73 | 0.36 | 1.50 | 0.375 |
| Cardiomyopathy | 1.23 | 0.52 | 2.92 | 0.623 |
| MAP | 1.01 | 1.00 | 1.02 | 0.078 |
| Cholesterol | 1.00 | 1.00 | 1.01 | 0.321 |
| Recipient age | 1.02 | 1.00 | 1.03 | 0.016 |
| Year of transplantation | 1.08 | 0.99 | 1.17 | 0.077 |
| Time of dialysis | 1.12 | 1.02 | 1.24 | 0.018 |
| CIT | 0.97 | 0.94 | 1.00 | 0.053 |
| Donor age | 0.99 | 0.98 | 1.00 | 0.130 |
| Sum of HLA mismatch | 0.95 | 0.83 | 1.09 | 0.451 |
| Hemoglobin | 1.00 | 0.85 | 1.18 | 0.989 |
| CNI usage | 0.06 | 0.04 | 0.10 | < 0.001 |
| Steroid usage | 11.29 | 6.59 | 19.33 | < 0.001 |