

**Single and Multivariable Models Adjusted with the Extended Set 1 of Confounders.**

**eTable 15 Single variable model with mAHEI and multivariable model adjusted with the extended set of confounders 1.**

Continuous independent variables	OR <sub>renal2vs1</sub>	OR <sub>renal3vs1</sub>	OR <sub>death2vs1</sub>	OR <sub>death3vs1</sub>	Median of tertile			P
					1	2	3	
mAHEI score	<b>0.96 (0.928-0.993)</b>	<b>0.872 (0.779-0.977)</b>	<b>0.859 (0.812-0.908)</b>	<b>0.601 (0.499-0.723)</b>	17.91	24.65	33.24	<b>&lt;0.0001</b>

Independent variables	OR <sub>renal2vs1</sub>	OR <sub>renal3vs1</sub>	OR <sub>death2vs1</sub>	OR <sub>death3vs1</sub>	Median of tertile			P
					1	2	3	
<b>Alcohol (drinks/week)</b>		<b>0.770 (0.632-0.937)</b>		<b>0.737 (0.57-0.953)</b>	0	0	5	<b>0.0129</b>
Animal proteins (g/kg/d)	0.972 (0.922-1.024)	0.925 (0.802-1.066)	1.014 (0.948-1.084)	1.039 (0.866-1.247)	0.27	0.47	0.81	0.4483
Plant proteins (g/kg/d)	1.011 (0.960-1.064)	1.03 (0.893-1.187)	0.988 (0.922-1.060)	0.968 (0.798-1.174)	0.14	0.2	0.3	0.8721
High-carbohydrate foods	1.023 (0.992-1.055)	1.122 (0.948-1.328)	1.026 (0.986-1.067)	1.163 (0.935-1.445)	2	9	21.46	0.1115
Deep fried food/snacks/fast food	0.896 (0.772-1.040)		1.036 (0.854-1.257)		no	yes (46.90%)		0.2385
<b>Fruits &amp; fruit juices</b>	<b>0.941 (0.891-0.995)</b>	<b>0.894 (0.808-0.990)</b>	<b>0.906 (0.845-0.972)</b>	<b>0.834 (0.733-0.948)</b>	4	9	17.82	<b>0.0037</b>
Vegetables	0.984 (0.926-1.045)	0.957 (0.814-1.124)	<b>0.905 (0.834-0.982)</b>	<b>0.766 (0.616-0.952)</b>	5	11	21	<b>0.0605</b>
24-hour urinary sodium (g)	0.975 (0.891-1.066)	0.975 (0.841-1.13)	0.909 (0.813-1.017)	0.953 (0.79-1.150)	3.46	4.89	6.4	0.1000
24-hour urinary potassium (g)	0.935 (0.872-1.003)	0.856 (0.727-1.007)	0.967 (0.881-1.061)	0.925 (0.744-1.149)	1.7	2.13	2.71	0.1808