

Single and Multivariable Models Adjusted with the Extended Set 1 of Confounders

eTable 18. Single variable model with mAHEI and multivariable model adjusted with the extended set of confounders 1.

Continuous independent variables	OR <sub>renal2vs1</sub>	OR <sub>renal3vs1</sub>	OR <sub>death2vs1</sub>	OR <sub>death3vs1</sub>	Median of tertile			p
					1	2	3	
<b>mAHEI score</b>	<b>0.899 (0.83-0.973)</b>	<b>0.787 (0.673-0.919)</b>	0.939 (0.821-1.073)	<b>0.659 (0.519-0.838)</b>	17.911	24.649	33.235	<0.0001

Independent variables	OR <sub>renal2vs1</sub>	OR <sub>renal3vs1</sub>	OR <sub>death2vs1</sub>	OR <sub>death3vs1</sub>	Median of tertile			p
					1	2	3	
<b>Alcohol (drinks/week)</b>		<b>0.809 (0.681-0.962)</b>		<b>0.74 (0.572-0.957)</b>	0	0	5	<b>0.0257</b>
Animal proteins (g/kg/d)	0.961 (0.916-1.007)	0.897 (0.789-1.019)	1.012 (0.946-1.082)	1.033 (0.86-1.24)	0.27	0.47	0.81	<b>0.0478</b>
Plant proteins (g/kg/d)	0.956 (0.913-1.002)	0.884 (0.776-1.007)	0.975 (0.910-1.046)	0.933 (0.77-1.132)	0.14	0.2	0.3	0.1388
High-carbohydrate foods	1.045 (1.016-1.075)	1.250 (1.075-1.454)	1.03 (0.991-1.072)	1.19 (0.957-1.48)	2	9	21.46	<b>0.0211</b>
Deep fried food/snacks/fast food	1.134 (0.995-1.293)		1.091 (0.900-1.324)		no	yes (46.90%)		0.1524
<b>Fruits &amp; fruit juices</b>	0.971 (0.923-1.021)	0.947 (0.863-1.040)	<b>0.911 (0.850-0.977)</b>	<b>0.841 (0.74-0.957)</b>	4	9	17.82	0.0700
<b>Vegetables</b>	0.978 (0.926-1.032)	0.941 (0.815-1.087)	<b>0.903 (0.832-0.980)</b>	<b>0.762 (0.613-0.947)</b>	5	11	21	<b>0.0332</b>
24-hour urinary sodium (g)	1.004 (0.926-1.089)	1.029 (0.901-1.176)	0.915 (0.818-1.023)	0.964 (0.799-1.163)	3.46	4.89	6.4	0.0795
<b>24-hour urinary potassium (g)</b>	<b>0.886 (0.832-0.943)</b>	<b>0.754 (0.652-0.871)</b>	0.953 (0.868-1.046)	0.893 (0.718-1.110)	1.7	2.13	2.71	0.0005