

## Mortality

**Supplemental table 3.** Multivariate Cox model with year of transplantation added, which was not included by the purposeful selection algorithm.

| <b>parameter</b>                       | <b>HR</b> | <b>95% CI</b> |      | <b>p-value</b> |
|--|-----------|---------------|------|----------------|
| <i>outcome mortality (n=106)</i>       |           |               |      |                |
| LA2D (per mm)                          | 1.06      | 1.03          | 1.08 | <0.001         |
| RV2D (per mm)                          | 0.96      | 0.90          | 1.02 | 0.14           |
| hemoglobin (per g/dl)                  | 0.81      | 0.71          | 0.92 | 0.001          |
| peripher vascular disease (yes vs. no) | 4.57      | 2.16          | 9.69 | <0.001         |
| immunosuppression (any vs. standard)   | 1.47      | 0.68          | 3.16 | 0.31           |
| immunosuppression (AZA vs MMF)         | 1.29      | 0.62          | 2.67 | 0.48           |
| CNI use (yes vs. no)                   | 1.51      | 0.51          | 4.43 | 0.43           |
| year of transplantation                | 0.97      | 0.91          | 1.04 | 0.43           |

HR, hazard ratio; 95% CI, 95% confidence interval; LA2D, left atrial diameter; RV2D, right ventricular diameter; AZA, azathioprine; MMF, mycophenolate; CNI, calcineurin inhibitor; standard immunosuppression regimen: steroid + MMF + CNI