

Multivariable Cox's proportional hazards model assessing the confounder-adjusted association of statin treatment on actual graft survival (graft failure and death with functioning graft counted as endpoints). \* Standard immunosuppression was steroids + MMF + CNI

	<b>HR</b>	<b>95% CI</b>	<b>p-value</b>
Statin treatment	0.68	0.55 – 0.84	<0.001
Age (per decade)	1.26	1.18 – 1.34	<0.001
Year of transplantation	0.96	0.93 – 0.99	0.014
Diabetes mellitus	1.24	1.00 – 1.54	0.049
Coronary heart disease	1.24	0.97 – 1.60	0.088
Peripheral arterial disease	1.33	1.03 – 1.73	0.030
Cardiomyopathy	1.21	0.91 – 1.59	0.185
Mean arterial pressure (per 10 mmHg)	1.02	1.00 – 1.04	0.037
No. of blood pressure drugs	1.08	1.02 – 1.15	0.009
Cholesterol (per 50 mg/dl)	0.89	0.82 – 0.96	0.003
Hemoglobin (g/dl)	0.83	0.80 – 0.86	<0.001
Immunosuppression (S + AZA + CNI vs. standard)*	1.20	0.90 – 1.61	0.214
Immunosuppression (steroid-free vs. standard)*	1.05	0.74 – 1.48	0.790
Immunosuppression (else vs. standard)*	1.56	1.20 – 2.02	<0.001