

Table 1 Demographic data of patients comparing the different CADI groups.

	No of drugs	CADI <2	CADI 2 - <4	CADI ≥ 4	p-value
No. of samples		52	21	9	
Donor (Living / deceased)		2/50	0/21	0/9	0.554
Donor creatinine (mg/dl)		0.81 (0.34)	0.80 (0.50)	0.97 (0.38)	0.290
Donor age (years)		41 (23)	59 (19)	63 (6)	<0.001
Recipient age (years)		52.1 (19.5)	61.8 (11.9)	65.4 (10.2)	0.006
Recipient weight (kg)		77 (18)	77.8 (11.5)	77.3 (21)	0.626
sex (male / female)		37 / 15	15 / 6	5 / 4	0.628
BCAR (%)		27	14	56	0.065
BCAR (days post TX)		13 (135)	8 (3)	14 (6)	0.205
PRA latest		2 (4)	0 (2)	0 (0)	0.101
CIT (hours)		12 (10)	12 (11)	9 (17)	0.821
DM		11	2	4	0.104*
Delayed graft function (no / yes)		50 / 2	18 / 3	8 / 1	0.160*
CMV (negative / positive)		26 / 10	10 / 6	3 / 4	0.316*
Pyelonephritis (negative / positive)		44 / 8	20 / 1	9 / 0	0.421*
Hyperlipidemia (number of drugs 0 / 1)		33 / 19	12 / 9	5 / 4	0.829
Cholesterol	0	199 (68)	220 (103)	208 (14)	0.785
	1	183 (76)	229 (121)	181 (45)	0.241
Hypertension (number of drugs 0 / 1 / 2 / 3 / 4 / 5)		4 / 11 / 16 / 9 / 10 / 2	3 / 1 / 2 / 5 / 6 / 4	0 / 1 / 2 / 2 / 4 / 0	0.134*
Mean arterial pressure (1 year post TX)	0	97 (17)	--	--	
	1	90 (18)	--	--	

	No of drugs	CADI <2	CADI 2 - <4	CADI ≥ 4	p-value
	2	93 (8)	90 (0)	98 (10)	0.263
	3	100 (13)	106 (12)	96 (0)	0.239
	4	94 (11)	92 (15)	95 (3)	0.487
	5	101 (12)	103 (10)	--	0.564
Immunosuppression: (Standard IS / steroid-free / else)		43/5 / 1	21/ 0 / 0	6/3 / 0	0.050*

Data represent median and interquartile range.

* Fisher test

Kruskal-Wallis test and Chi-square test were used for computation of p-values

nd no data; na not applicable, standard IS ... steroids+MMF+CNI